

CLIMATE CHANGE PROJECTIONS FOR INDIVIDUAL TREE SPECIES NORTHERN SUPERIOR UPLANDS (ECOLOGICAL SECTION 212L)



This region's forests will be affected by a changing climate and other stressors during this century. A team of managers and researchers created an assessment that describes the vulnerability of forests in the region ([Handler et al. 2014](#)). This report includes information on observed and future climate trends, and also summarizes key vulnerabilities for forested natural communities. The Landscape Change Research Group recently updated the Climate Change Tree Atlas, and this handout summarizes that information.

Full Tree Atlas results are available online at www.fs.fed.us/nrs/atlas/. Two climate scenarios are presented to “bracket” a range of possible futures. These future climate projections (2070 to 2099) provide information about how individual tree species may respond to a changing climate. Results for “low” and “high” emissions scenarios can be compared on the reverse side of this handout.

The updated Tree Atlas presents additional information helpful to interpret tree species changes:

- **Suitable habitat** - calculated based on 39 variables that explain where optimum conditions exist for a species, including soils, landforms, and climate variables.
- **Adaptability** - based on life-history traits that might increase or decrease tolerance of expected changes, such as the ability to withstand different forms of disturbance.
- **Capability** - a rating of the species' ability to cope or persist with climate change in this region based on suitable habitat change (statistical modeling), adaptability (literature review and expert opinion), and abundance (FIA data). The capability rating is modified by abundance information; ratings are downgraded for rare species and upgraded for abundant species.
- **Migration Potential Model** - when combined with habitat suitability, an estimate of a species' colonization likelihood for new habitats. This rating can be helpful for assisted migration or focused management (see the table section: “New Habitat with Migration Potential”).

Remember that models are just tools, and they're not perfect. Model projections can't account for all factors that influence future species success. If a species is rare or confined to a small area, model results may be less reliable. These factors, and others, could cause a particular species to perform better or worse than a model projects. Human choices will also continue to influence forest distribution, especially for tree species that are projected to increase. Planting programs may assist the movement of future-adapted species, but this will depend on management decisions. Despite these limits, models provide useful information about future expectations. It's perhaps best to think of these projections as indicators of possibility and potential change.

SOURCE: This handout summarizes the full model results for the Northern Superior Uplands (Ecological Section 212L), available at www.fs.fed.us/nrs/atlas/combined/resources/summaries. More information on vulnerability and adaptation in the region can be found at www.forestadaptation.org/northwoods. A full description of the models and variables are provided in Iverson et al. 2019 (www.nrs.fs.fed.us/pubs/57857 and www.nrs.fs.fed.us/pubs/59105) and Peters et al. 2019 (www.nrs.fs.fed.us/pubs/58353).

CLIMATE CHANGE CAPABILITY

POOR CAPABILITY

American mountain-ash	Pin cherry
Balsam fir	Serviceberry
Balsam poplar	White spruce
Black willow	

FAIR CAPABILITY

Black cherry	Quaking aspen
Black spruce	Red pine
Mountain maple	Tamarack (native)
Paper birch	

GOOD CAPABILITY

American basswood	Jack pine
American elm	Northern pin oak
Bigtooth aspen	Northern red oak
Boxelder	Red maple
Bur oak	Slippery elm
Eastern white pine	Sugar maple
Green ash	Yellow birch
Ironwood	

MIXED RESULTS

Black ash	Silver maple
Northern white-cedar	

NEW HABITAT WITH MIGRATION POTENTIAL

American beech	Eastern redbud
American hornbeam	Eastern redcedar
Ashe juniper	Hackberry
Bigleaf magnolia	Honeylocust
Bitternut hickory	Live oak
Black locust	Scarlet oak
Black oak	Shagbark hickory
Black walnut	Swamp white oak
Cedar elm	Sweet birch
Chestnut oak	Sweetgum
Chinkapin oak	Sycamore
Eastern cottonwood	White ash
Eastern hemlock	White oak



ADAPTABILITY: Life-history factors, such as the ability to respond favorably to disturbance, that are not included in the Tree Atlas model and may make a species more or less able to adapt to future stressors.

- + **HIGH** *Species may perform better than modeled*
- **MEDIUM**
- **LOW** *Species may perform worse than modeled*

HABITAT CHANGE: Projected change in suitable habitat between current and potential future conditions.

- ▲ **INCREASE** *Projected increase of >20% by 2100*
- **NO CHANGE** *Projected change of <20% by 2100*
- ▼ **DECREASE** *Projected decrease of >20% by 2100*
- ★ **NEW HABITAT** *Tree Atlas projects new habitat for species not currently present*

ABUNDANCE: Based on Forest Inventory Analysis (FIA) summed Importance Value data, calibrated to a standard geographic area.

- + **ABUNDANT**
- **COMMON**
- **RARE**

CAPABILITY: An overall rating that describes a species' ability to cope or persist with climate change based on suitable habitat change class (statistical modeling), adaptability (literature review and expert opinion), and abundance within this region.

- ▲ **GOOD** *Increasing suitable habitat, medium or high adaptability, and common or abundant*
- **FAIR** *Mixed combinations, such as a rare species with increasing suitable habitat and medium adaptability.*
- ▼ **POOR** *Decreasing suitable habitat, medium or low adaptability, and uncommon or rare*

SPECIES	LOW CLIMATE CHANGE (RCP 4.5)				HIGH CLIMATE CHANGE (RCP 8.5)			
	HABITAT		CAPABILITY		HABITAT		CAPABILITY	
	ADAPT	ABUN	CHANGE	CHANGE	CHANGE	CHANGE	CHANGE	CHANGE
American basswood	•	-	▲	△	▲	△		
American beech	•		★		★			
American elm	•	-	▲	△	▲	△		
American hornbeam*	•		★		★			
American mountain-ash*	-	-	▼	▽	▼	▽		
Ashe juniper	•		★		★			
Balsam fir	-	+	▼	▽	▼	▽		
Balsam poplar	•	•	▼	▽	▼	▽		
Bigleaf magnolia*	•				★			
Bigtooth aspen	•	•	▲	△	▲	△		
Bitternut hickory*	+		★		★			
Black ash	-	+	▲	△	●	○		
Black cherry	-	-	▲	○	▲	○		
Black locust*	•		★		★			
Black oak	•		★		★			
Black spruce	•	+	▼	○	▼	○		
Black walnut*	•		★		★			
Black willow*	-	-	▼	▽	●	▽		
Boxelder*	+	-	▲	△	▲	△		
Bur oak	+	-	▲	△	▲	△		
Cedar elm	-		★		★			
Chestnut oak	+		★		★			
Chinkapin oak	•				★			
Eastern cottonwood*	•		★		★			
Eastern hemlock	-		★		★			
Eastern redbud*	•				★			
Eastern redcedar	•		★		★			
Eastern white pine	-	•	▲	△	▲	△		
Green ash*	•	-	▲	△	▲	△		
Hackberry	+		★		★			
Honeylocust*	+				★			
Ironwood*	+	-	▲	△	▲	△		
Jack pine	+	+	▼	△	▼	△		
Live oak	•		★		★			
Mountain maple*	+	•	▼	○	▼	○		
Northern pin oak	+	-	▲	△	▲	△		
Northern red oak	+	-	▲	△	▲	△		
Northern white-cedar	•	+	▼	○	●	△		
Paper birch	•	+	▼	○	▼	○		
Pin cherry*	•	-	▼	▽	▼	▽		
Quaking aspen	•	+	▼	○	▼	○		
Red maple	+	+	▲	△	▲	△		
Red pine	-	•	▲	○	▲	○		
Scarlet oak	•		★		★			
Serviceberry*	•	-	▼	▽	▼	▽		
Shagbark hickory	•		★		★			
Silver maple*	+	-	●	○	▲	△		
Slippery elm*	•	-	▲	△	▲	△		
Sugar maple	+	•	▲	△	▲	△		
Swamp white oak*	•		★		★			
Sweet birch	-		★		★			
Sweetgum	•				★			
Sycamore*	•		★		★			
Tamarack (native)	-	•	▲	○	▲	○		
White ash	-		★		★			
White oak	+		★		★			
White spruce	•	•	▼	▽	▼	▽		
Yellow birch	•	•	▲	△	▲	△		

*Species with low model reliability based on five statistical metrics of the habitat models that affect change class. See maps and tables for more information (www.fs.fed.us/nrs/atlas/combined/resources/summaries).